

Arctic Report Card 2016

Persistent warming trend and loss of sea ice are triggering extensive Arctic changes

2016 Headlines

Executive Summary

2016 Addendum

Contacts

Vital Signs

Surface Air Temperature

Terrestrial Snow Cover

Greenland Ice Sheet

Sea Ice

Sea Surface Temperature

Arctic Ocean Primary Productivity

Tundra Greenness

Indicators

Ocean Acidification

Terrestrial Carbon Cycle

Shrews and Their Parasites: Small Species Indicate Big Changes

Frostbites

Arctic Change – So What?:

Linkages and Impacts

Faster Glaciers and the Search for Faster Science

More Information

About Arctic Report Card 2016

Authors and Affiliations

References

2016 Headlines

Persistent warming trend and loss of sea ice are triggering extensive Arctic changes.

Observations in 2016 showed a continuation of long-term Arctic warming trends which reveals the interdependency of physical and biological Arctic systems, contributing to a growing recognition that the Arctic is an integral part of the globe, and increasing the need for comprehensive communication of Arctic change to diverse user audiences.

Highlights

- The average **surface air temperature** for the year ending September 2016 is by far the highest since 1900, and new monthly record highs were recorded for January, February, October and November 2016.
- After only modest changes from 2013-2015, **minimum sea ice extent** at the end of summer 2016 tied with 2007 for the second lowest in the satellite record, which started in 1979.
- **Spring snow cover extent** in the North American Arctic was the lowest in the satellite record, which started in 1967.
- In 37 years of **Greenland ice sheet** observations, only one year had earlier onset of spring melting than 2016.
- The Arctic Ocean is especially prone to **ocean acidification**, due to water temperatures that are colder than those further south. The short Arctic food chain leaves Arctic marine ecosystems vulnerable to ocean acidification events.
- Thawing permafrost releases **carbon** into the atmosphere, whereas greening tundra absorbs atmospheric carbon. Overall, tundra is presently releasing net carbon into the atmosphere.
- Small Arctic mammals, such as **shrews**, and their parasites, serve as indicators for present and historical environmental variability. Newly acquired parasites indicate northward shifts of sub-Arctic species and increases in Arctic biodiversity.

Video



<http://www.arctic.noaa.gov/Report-Card>